

CONCLUSION

CAS is as technically feasible, safe, and durable in anatomically high-risk patients as in medically high-risk patients, with similar rates of periprocedural stroke, death, and late restenosis. However, patients with radiation-induced stenosis appear to be at an increased risk for restenosis.

AUTHOR CONTRIBUTIONS

Conception and design: SS, JP, AR
Analysis and interpretation: SS, JP
Data collection: SS, AR, CS
Writing the article: SS, JP
Critical revision of the article: SS, JP
Final approval of the article: SS, JP
Statistical analysis: SS, CS
Obtained funding: JP, RD, RS
Overall responsibility: JP

REFERENCES

- Gagne PJ, Riles TS, Jacobowitz GR, Lamparello PJ, Giangola G, Adelman MA, et al. Long-term follow-up of patients undergoing reoperation for recurrent carotid artery disease. *J Vasc Surg* 1993;18:991-8; discussion 999-1001.
- Hassen-Khodja R, Sala F, Declémy S, Lagrange JL, Bouillane PJ, Batt M. Surgical management of atherosclerotic carotid artery stenosis after cervical radiation therapy. *Ann Vasc Surg* 2000;14:608-11.
- AbuRahma AF, Jennings TG, Wulu JT, Tarakji L, Robinson PA. Redo carotid endarterectomy versus primary carotid endarterectomy. *Stroke* 2001;32:2787-92.
- Mansour MA, Kang SS, Baker WH, Watson WC, Littooy FN, Labropoulos N, et al. Carotid endarterectomy for recurrent stenosis. *J Vasc Surg* 1997;25:877-83.
- Diethrich EB, Ndiaye M, Reid DB. Stenting in the carotid artery: initial experience in 110 patients. *J Endovasc Surg* 1996;3:42-62.
- Yadav JS, Roubin GS, Iyer S, Vitek J, King P, Jordan WD, et al. Elective stenting of the extracranial carotid arteries. *Circulation* 1997;95:376-81.
- Bergeron P, Becquemin JP, Jausseran JM, Biasi G, Cardon JM, Castellani L, et al. Percutaneous stenting of the internal carotid artery: the European CAST I study. Carotid artery stent trial. *J Endovasc Surg* 1999;6:155-9.
- Yadav JS, Wholey MH, Kuntz RE, Fayad P, Katzen BT, Mishkel GJ, et al. Protected carotid-artery stenting versus endarterectomy in high-risk patients. *N Engl J Med* 2004;351:1493-501.
- Fitzpatrick CM, Chiou AC, DeCaprio JD, Kashyap VS. Carotid revascularization in the presence of contralateral carotid artery occlusion is safe and durable. *Mil Med* 2005;170:1069-74.
- Domenig C, Hamdan AD, Belfield AK, Campbell DR, Skillman JJ, LoGerfo FW, et al. Recurrent stenosis and contralateral occlusion: high-risk situations in carotid endarterectomy? *Ann Vasc Surg* 2003;17:622-8.
- AbuRahma AF, Snodgrass KR, Robinson PA, Wood DJ, Meek RB, Patton DJ. Safety and durability of redo carotid endarterectomy for recurrent carotid artery stenosis. *Am J Surg* 1994;168:175-8.
- Elerding SC, Fernandez RN, Grotta JC, Lindberg RD, Causay LC, McMurtrey MJ. Carotid artery disease following external cervical irradiation. *Ann Surg* 1981;194:609-15.
- Moritz MW, Higgins RF, Jacobs JR. Duplex imaging and incidence of carotid radiation injury after high-dose radiotherapy for tumors of the head and neck. *Arch Surg* 1990;125:1181-3.
- Atkinson JL, Sundt TM Jr, Dale AJ, Cascino TL, Nichols DA. Radiation-associated atheromatous disease of the cervical carotid artery: report of seven cases and review of the literature. *Neurosurgery* 1989;24:171-8.
- Kashyap VS, Moore WS, Quinones-Baldrich WJ. Carotid artery repair for radiation-associated atherosclerosis is a safe and durable procedure. *J Vasc Surg* 1999;29:90-6; discussion 97-9.
- Ferguson GG, Eliasziw M, Barr HW, Clagett GP, Barnes RW, Wallace MC, et al. The North American Symptomatic Carotid Endarterectomy Trial: surgical results in 1415 patients. *Stroke* 1999;30:1751-8.
- Melliere D, Becquemin JP, Berrahal D, Desgranges P, Cavillon A. Management of radiation-induced occlusive arterial disease: a reassessment. *J Cardiovasc Surg (Torino)* 1997;38:261-9.
- Teitelbaum GP, Lefkowitz MA, Giannotta SL. Carotid angioplasty and stenting in high-risk patients. *Surg Neurol* 1998;50:300-11; discussion 311-2.
- Cheng SW, Wu LL, Ting AC, Lau H, Lam LK, Wei WI. Irradiation-induced extracranial carotid stenosis in patients with head and neck malignancies. *Am J Surg* 1999;178:323-8.
- Eskandari MK, Brown KE, Kibbe MR, Morasch MD, Matsumura JS, Pearce WH. Restenosis after carotid stent placement in patients with previous neck irradiation or endarterectomy. *J Vasc Interv Radiol* 2007;18:1368-74.
- Zahn R, Mark B, Niedermaier N, Zeymer U, Limbourg P, Ischinger T, et al. Embolic protection devices for carotid artery stenting: better results than stenting without protection? *Eur Heart J* 2004;25:1550-8.
- Grube E, Gerckens U, Yeung AC, Rowold S, Kirchhof N, Sedgewick J, et al. Prevention of distal embolization during coronary angioplasty in saphenous vein grafts and native vessels using porous filter protection. *Circulation* 2001;104:2436-41.
- Terada T, Yamaga H, Tsumoto T, Masuo O, Itakura T. Use of an embolic protection system during endovascular recanalization of a totally occluded cervical internal carotid artery at the chronic stage. Case report. *J Neurosurg* 2005;102:558-64.
- Protack CD, Bakken AM, Saad WE, Illig KA, Waldman DL, Davies MG. Radiation arteritis: a contraindication to carotid stenting? *J Vasc Surg* 2007;45:110-7.

Submitted Feb 22, 2009; accepted Apr 24, 2009.

DISCUSSION

Dr Peter Lin (Houston, Tex). Dr Shin and her colleagues from the Eastern Virginia Medical School should be congratulated for an excellent clinical study in which they analyzed the clinical outcome of carotid stenting in patients who have high-risk medical comorbidities vs those with anatomically high-risk lesions. The authors reported similar clinical outcome in terms of stroke and mortality between the two groups. However, patients with radiation-induced stenosis had a much higher risk of restenosis compared to the medically high-risk cohorts. I have three questions for the author.

My first question relates to the ultrasound surveillance. The higher risks of restenosis in patients with radiation-induced carotid

lesions, in my view, warrants a more vigilant or frequent ultrasound surveillance protocol since the need for reintervention is undoubtedly higher. Can you share with us your view whether a more vigilant surveillance protocol is necessary in this particular patient cohort? If so, what should be the ideal surveillance protocol in these patients?

Second, in your study, a variety of carotid stents including the Smart, Precise, and Acculink stents were used. There have been studies in the recent literature to suggest these nitinol stents, which are considered a open-cell stent, have lower patency compared to closed-cell stents such as Wallstent. This is a finding that we have also validated in our own clinical experience. This finding has also

led some researchers to advocate the notion that different carotid stent should be applied based on patients' anatomical factors. Can you share with us with your opinion regarding the notion that different devices (ie, open- vs closed-cell stent) should be applied to different patient cohorts based on the higher risk of restenosis?

My last question has to do with the ultrasound criteria which you based to determine the degree of restenosis. In our practice, we have found the velocity criteria used to determine carotid stenting-related restenosis varies based on the stent used. In other words, we have noted that the same velocity criteria used for a nitinol stent does not apply to the Wallstent. We also learned that same velocity criteria for de novo carotid lesion do not apply to lesion of carotid stent-related restenosis. How did you derive your velocity criteria in your study? Have you found a difference in velocity criteria based on the stents used in your study?

I'd like to thank the authors for providing me with a well-written manuscript well ahead of time. I'd also want to thank the association for the opportunity to discuss this paper.

Dr Susanna H. Shin. Thank you for your comments and your questions. Your first question is on the surveillance of our patients. Routinely, our patients undergo a duplex ultrasound within 24 hours of their procedure, then at 1 month and 6

months, and annually thereafter. In our study, we had patients in the radiation cohort who developed restenosis at 7.5 months, 9.1 months, 13 months, and at 58 months. Currently, we do not have a different protocol for following up these patients more often, perhaps that would lead us to detecting restenoses earlier and potentially provide for them to have reintervention sooner.

Your second question was based on the type of stent. The majority of our patients were part of a trial. Obviously, the stent choice was predetermined, and therefore dictated by that trial and the majority of our patients received open-cell stents. Future directions would be potentially to use closed-cell stents, which might be better in this specific cohort of radiation-induced stenosis, and perhaps another option would be using drug-eluting stents or covered stents.

For stenosis criteria, we don't use, at this time, different velocity criteria for different stents. We currently use the established criteria set forth by Dr AbuRhamah and currently we are looking at our own duplex velocity data and comparing them to our angiograms to see how they correlate and potentially come up with new criteria for those stents.

Access to *Journal of Vascular Surgery Online* is reserved for print subscribers!

Full-text access to *Journal of Vascular Surgery Online* is available for all print subscribers. To activate your individual online subscription, please visit *Journal of Vascular Surgery Online*, point your browser to <http://www.jvascsurg.org>, follow the prompts to **activate your online access**, and follow the instructions. To activate your account, you will need your subscriber account number, which you can find on your mailing label (*note*: the number of digits in your subscriber account number varies from 6 to 10). See the example below in which the subscriber account number has been circled:

Sample mailing label

This is your subscription
account number

*****3-DIGIT 001	
SJ P1	
FEB00 J024 C: 1	(1234567-89) U 05/00 Q: 1
J. H. DOE, MD	
531 MAIN ST	
CENTER CITY, NY 10001-001	

Personal subscriptions to *Journal of Vascular Surgery Online* are for individual use only and may not be transferred. Use of *Journal of Vascular Surgery Online* is subject to agreement to the terms and conditions as indicated online.